

Contact: Christian Basi  
[BasiC@missouri.edu](mailto:BasiC@missouri.edu)  
573-882-4430  
[University of Missouri-Columbia](#)


## MU scientists study how to improve pesticide efficiency

COLUMBIA, Mo. – In 2007, a controversial pesticide was approved by the Food and Drug Administration for use on fruit and vegetable crops, mainly in California and Florida. Farm workers and scientists protested the approval of the pesticide because its active ingredient, methyl iodide, is a known carcinogen. Now, MU researchers are studying the molecular structure of the pesticide to determine if the product could be made more efficient and safer for those living near, and working in, treated fields.

Methyl iodide is the active ingredient used in a pesticide known commercially as Midas. Midas is a mix of methyl iodide and chloropicrin, a rat poison, and is used primarily on the fields that will grow strawberries, tomatoes and bell peppers. In a new study published this month in the *Journal of Agricultural and Food Chemistry*, MU researchers studied why the manufacturer blended the chemicals to determine if a different chemical combination might be possible.


"We found that the two chemicals, methyl iodide and chloropicrin, are mixed to slow the release of methyl iodide and increase its effectiveness," said Rainer Glaser, professor of chemistry in the MU College of Arts & Science. "However, we believe that a different chemical mix could further slow the release of methyl iodide and allow farmers to use less of the pesticide, which would make the area safer for workers and the public."



 **IMAGE:** Rainer Glaser, professor of chemistry in the MU College of Arts & Science, studied the molecular structure of a pesticide to determine if the product could be made more efficient...

[Click here for more information.](#)



 **IMAGE:** Kaitlan Prugger, a co-author and undergraduate researcher, studied Midas, a commercially available pesticide, to better understand its molecular structure and how it could be made safer and more efficient.

[Click here for more information.](#)

Methyl iodide is a fumigant, meaning that it fills an air space with gas, suffocating and poisoning the pests within the airspace. Farm workers dressed in protective suits apply Midas in liquid form to fields 10 to 14 days prior to planting and cover future crop rows with plastic sheeting. During that time period, the pesticide is released in gas form, suffocating pests in the top layer of the soil. However, much of the fumigant is useless as it is lost into the atmosphere upon application and during the release period due to ventilation of the area. All of the pesticide is gone before the area is planted.

"Farmers use 200 to 300 pounds of Midas per acre and nearly 80 percent of the pesticide is not effective in killing pests," said Kaitlan Prugger, a co-author and undergraduate researcher. "Gas lost to the atmosphere could pose risks to farm workers and nearby communities. Even a small improvement in effectiveness achieved through a change in the chemical blend could greatly reduce the amount of pesticide used per acre."

The use of methyl iodide is a consequence of the Montreal Protocol on Substances that Deplete the Ozone Layer. Prior to the protocol, farmers used methyl bromide to fumigate strawberry and tomato fields. However, methyl bromide was found to deplete the ozone and its use was phased out completely in 2005.

###



*AAAS and EurekAlert! are not responsible for the accuracy of news releases posted to EurekAlert! by contributing institutions or for the use of any information through the EurekAlert! system.*

---

[HOME](#) [DISCLAIMER](#) [PRIVACY POLICY](#) [TERMS & CONDITIONS](#) [CONTACT US](#) [TOP](#)  
Copyright ©2012 by AAAS, the science society.